
TA-274-0116

Aedes (Aedimorphus) vexans (Meigen), WRBU specimen AEvex, Character descriptions: Carpenter and LaCasse, 1955: 263



Aedes (Aedimorphus) vexans (Meigen), WRBU specimen AEvex, Character descriptions: Carpenter and LaCasse, 1955: 263










Bionomics: Immatures are found in unshaded fresh water flood pools in secondary scrub, but have also been collected in ditches, swamps, rice fields, and elephant foot prints. Habitats usually have little aquatic vegetation or algae. Females are night biters and readily feed on man and cattle (Reinert, 1973).

Medical Importance: Capable of transmitting eastern equine encephalitis virus (EEE), western equine encephalitis virus (WEE), SLE, West Nile Virus (WNV) (Turell *et al.*, 2005:60). It is also a vector of dog heartworm (Reinert, 1973). Troublesome biter. Flight range up to 10 miles.



HindLeg. Hind tarsi dark, with narrow basal white rings on all segments; front and middle tarsi similarly marked, but with white rings on segments 4 and 5 greatly reduced or absent.

WALTER REED BIOSYSTEMATICS UNIT

Army Public Health Center (Provisional), Entomological Sciences Program / COM 410-436-3613 / DSN 312-584-3613 / Website: http://phc.amedd.army.mil / Approved for Public Release, Distribution Unlimited